

Justo Galeas  
 EDG500  
 04/31/23

**Descriptives**

Reported pain level

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum
					Lower Bound	Upper Bound	
Low dosage	3	7.0000	1.00000	.57735	4.5159	9.4841	6.00
Moderate dosage	3	5.6667	2.08167	1.20185	.4955	10.8378	4.00
High dosage	3	2.0000	1.00000	.57735	-.4841	4.4841	1.00
Total	9	4.8889	2.57121	.85707	2.9125	6.8653	1.00

**ANOVA**

Reported pain level

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	40.222	2	20.111	9.526	.014
Within Groups	12.667	6	2.111		
Total	52.889	8			

**Descriptives**

Hours of internet usage weekly

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
low ses	4	10.0000	1.82574	.91287	7.0948	12.9052	8.00	12.00
middle ses	4	12.2500	1.70783	.85391	9.5325	14.9675	10.00	14.00
high ses	4	12.0000	2.16025	1.08012	8.5626	15.4374	10.00	15.00
Total	12	11.4167	2.02073	.58333	10.1328	12.7006	8.00	15.00

**ANOVA**

Hours of internet usage weekly

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	12.167	2	6.083	1.672	.241
Within Groups	32.750	9	3.639		
Total	44.917	11			

- A. The low-ses group mean is 10.00.
- B. The middle-ses group mean is 12.25.
- C. The high-ses group mean is 12.00.
- D. The value of F is 1.67.
- E. The associated probability is .241.
- F. .241 is greater than .05.
- G. The mean for the high-ses group is 12.00 (sd = 2.16), for the middle-ses group is 12.25 (sd = 1.71), and for the low-ses group is 10.00 (sd = 1.82). The differences among the means are not statistically significant at the .05 level (F = 1.67, p = .241).