

DEMONSTRATION CHECKLIST FOR ADMINISTRATION AND DOCUMENTATION OF MEDICATIONS

TABLET/CAPSULE, LOZENGE MEDICATIONS	RATIONALE						
<p>7/19 1. Washed hands.</p> <p>7/19 2. Unlocked medication cabinet.</p> <p>7/19 3. Checked individual's monthly medication sheet to determine medications to be administered.</p> <p>7/19 4. Assembled equipment necessary for administration.</p> <p>7/19 5. Named 2 sources to find the purpose, side effects, and any warnings for the medication.</p> <p>7/19 6. Checked for allergies to medication.</p> <p>7/19 7. Removed medication from individual's supply and compared the medication label against individual's medication sheet for:</p> <table style="margin-left: 40px; border: none;"> <tr> <td>7/19 Right Individual</td> <td>7/19 Right Medication</td> </tr> <tr> <td>7/19 Right Date</td> <td>7/19 Right Time</td> </tr> <tr> <td>7/19 Right Route</td> <td>7/19 Right Dose</td> </tr> </table> <p>7/19 8. Checked expiration date.</p> <p>7/19 9. Identified what to do if medication label does not match medication sheet.</p> <p>7/19 10. Compared medication label against individual's medication sheet for the 2nd time.</p> <p>7/19 11. For medications in a bottle: poured correct number of tablets/capsules into the lid of the medication container and transferred them into a medication cup. For medications in a 'bubble pack': started at the highest number, pushed the correct dosage into a medication cup, and wrote the date and their initials on the card next to the dosage(s) popped out. For lozenges: unwrapped the lozenge and transferred it into a medication cup.</p> <p>7/19 12. Compared medication label against individual's medication sheet for the 3rd time.</p> <p>7/19 13. Identified individual prior to administration of medication.</p> <p>7/19 14. Explained to individual what is to be done.</p> <p>7/19 15. Administered correct dose of medication by instructing individual to swallow meds (offered min. 4 oz. water). If the medication is in lozenge form, instructed individual not to chew or swallow; the lozenge needs to dissolve in their mouth.</p> <p>7/19 16. For swallowed medication: remained with individual until medication was swallowed. For lozenges: remained in same area of the individual until the lozenge was completely dissolved. Checked to ensure individual did not chew or swallow the lozenge.</p> <p>7/19 17. Returned medication to locked area.</p> <p>7/19 18. Disposed of used supplies.</p> <p>7/19 19. Washed hands.</p> <p>7/19 20. Charted medication administered correctly.</p>	7/19 Right Individual	7/19 Right Medication	7/19 Right Date	7/19 Right Time	7/19 Right Route	7/19 Right Dose	<p>1. To prevent the spread of disease.</p> <p>2. To ensure individual safety, medications are kept locked.</p> <p>3. To review correct medication orders.</p> <p>4. To be organized.</p> <p>5. To be informed about the medication being given.</p> <p>6. To avoid giving medication that a person is allergic to.</p> <p>7. To prevent medication errors.</p> <p>8. To avoid administering ineffective medication.</p> <p>9. To know what steps to take.</p> <p>10. To verify accuracy of 1st check.</p> <p>11. To follow correct and sanitary procedures for medication administration.</p> <p>12. To verify accuracy of 2nd check.</p> <p>13. To avoid giving medication to the wrong individual.</p> <p>14. To ensure individual understands medication procedure.</p> <p>15. To administer medication as ordered.</p> <p>16. To ensure entire dose is taken.</p> <p>17. To ensure individual safety, medications are kept locked.</p> <p>18. To clean the area.</p> <p>19. To prevent the spread of disease.</p> <p>20. To follow policy and procedure on medication administration and documentation.</p>
7/19 Right Individual	7/19 Right Medication						
7/19 Right Date	7/19 Right Time						
7/19 Right Route	7/19 Right Dose						